arXiv:1612.00288v1 [physics.gen-ph] 27 Nov 2016

A testable prediction from entropic gravity

Mehmet Süzen

(Dated: December 2, 2016)

I have shown conceptually that quantum state has a direct relationship to gravitational constant due to entropic force posed by Verlinde's argument and part of the Newton-Schrödinger equation (N-S) in the context of gravity induced collapse of the wavefunction via Diósi-Penrose proposal. This direct relationship can be used to measure gravitational constant using state-of-the-art mater-wave interforemetry to test the entropic gravity argument.

INTRODUCTION

Recently, Verlinde has put forward an argument regarding entropic force induced gravity [1]. On the other hand, Newton-Schrödinger equation is developed to couple gravity with the collapse of a wavefunction, Penrose-Diósi [2–5]. Wavefunction collapse implies that when quantity of mass approaching to macroscopic scales, the wavefunction tends to localize in position [6]. Here we aim at using these two approaches to drive a consistent relationship between quantum state of atomic system and the gravitational constant.

ENTROPIC WAVEFUNCTION

Unruh radiation is observed for an accelerated particle with acceleration a [7] with temperature T,

$$T = \frac{1}{2\pi}\frac{\hbar a}{k_B c} \tag{1}$$

where, $\hbar$ is the Planck constant, k_B is Boltzmann constant and c is the speed of light. If we consider this particle placed in front of a membrane, holographic screen [1], its distance and entropy associated with the information on the membrane will read as follows,

$$\Delta x = \frac{\hbar}{mc}, \tag{2}$$

$$\Delta S = 2\pi k_B. \tag{3}$$

This originates from Bekenstein's thought experiment [1]. The force F excerted by the membrane on the particle reads,

$$F = T\frac{\Delta S}{\Delta x}. \tag{4}$$

We can plug Eqs. (1), (2) and (3) into Eq. (4). This will lead to Newton's Law, with scalar field, particle's own graviational field, $\phi(x)$,

$$F = ma = -\nabla\phi(x) = -\frac{d\phi(x)}{dx}. \tag{5}$$

If we consider particle as a quantum mechanical object, moving in its own gravitational field, the Schrödinger-Newton equation (S-N), are the pair of coupled nonlinear partial differential equations [8],

$$\nabla^2\phi(x) = 4\pi G m|\psi|^2, \tag{6}$$

$$-\frac{\hbar^2}{2m}\nabla^2\psi - m\phi\psi = \tilde{E}\psi. \tag{7}$$

where $\phi(x)$ is the scalar field, m is the mass, ψ is the wavefunction, G is the gravitational constant and $\tilde{E}$ is the energy. Argument due to Penrose [4] is that multiple quantum states reduce to one of the states in finite time because of the significant mass displacement.

Re-writing the first part of S-N with the entropic force, Eq.(6) with Eqs.(4) and (5) in one dimension,

$$\frac{d}{dx}(F) = 4\pi G m|\psi|^2, \tag{8}$$

$$\frac{d}{dx}(-\frac{d\phi(x)}{dx}) = 4\pi G m|\psi|^2, \tag{9}$$

$$-\frac{d}{dx}(T\frac{\Delta S}{\Delta x}) = 4\pi G m|\psi|^2. \tag{10}$$

Integrating both sides on Eq.(10) and recall Δx

$$-T\frac{mc\Delta S}{\hbar} + C = 4\pi G m\int|\psi|^2 dx. \tag{11}$$

, If we write the entropy with von Neumann entropy *without the trace for collapsed state*,

$$\Delta S = -k_B\hat{\rho}\ln\hat{\rho}, \tag{12}$$

where $\hat{\rho}$ is the density matrix, it's proportional to collapsed density,

$$\hat{\rho} \sim \int|\psi|^2 dx. \tag{13}$$

Using Eq.(12) and (13) into Eq.(11), with T and Δx,

$$\frac{-1}{2\pi}\frac{\hbar a}{k_B c}\frac{mck_B ln\hat{\rho}}{\hbar} \sim 4\pi G m\hat{\rho}, \tag{14}$$

$$\hat{\rho} \sim exp(-8\pi^2 G/a). \tag{15}$$

CONCLUSION

The proposed relationship at Eq. (15) could be used in measuring G using atom interforemeter [9] to varify entropic gravity argument. However, mass of the atomic system should be $10^9 u$ to go into S-N collapse [6], this

might be an experimental challange [6], and density matrix of the atomic system should be computable from the first-principles.

[1] Erik Verlinde. On the origin of gravity and the laws of newton. *Journal of High Energy Physics*, 2011(4):1–27, 2011.

[2] Lajos Diosi. A universal master equation for the gravitational violation of quantum mechanics. *Physics letters A*, 120(8):377–381, 1987.

[3] LAJOS Diosi. Models for universal reduction of macroscopic quantum fluctuations. *Physical Review A*, 40(3):1165, 1989.

[4] Roger Penrose. On gravity's role in quantum state reduction. *General relativity and gravitation*, 28(5):581–600, 1996.

[5] Lajos Diósi. Notes on certain newton gravity mechanisms of wavefunction localization and decoherence. *Journal of Physics A: Mathematical and Theoretical*, 40(12):2989, 2007.

[6] James R van Meter. Schrödinger-newton collapse of the wavefunction. *Classical and Quantum Gravity*, 28(21):215013, 2011.

[7] William G Unruh. Notes on black-hole evaporation. *Physical Review D*, 14(4):870, 1976.

[8] Richard Harrison, Irene Moroz, and KP Tod. A numerical study of the schrödinger–newton equations. *Nonlinearity*, 16(1):101, 2003.

[9] Jeffrey B Fixler, GT Foster, JM McGuirk, and MA Kasevich. Atom interferometer measurement of the newtonian constant of gravity. *Science*, 315(5808):74–77, 2007.